

Blazar Heating – The Rosetta Stone for Structure Formation?

Christoph Pfrommer¹

in collaboration with

Avery E. Broderick², Phil Chang², Ewald Puchwein¹, Volker Springel¹

¹Heidelberg Institute for Theoretical Studies, Germany

²Canadian Institute for Theoretical Astrophysics, Canada

Sep 13, 2011 / Stockholm University/OKC



Outline

- 1 Physics of blazar heating
 - TeV emission from blazars
 - Propagation of TeV photons
 - Plasma instabilities
- 2 The intergalactic medium
 - Properties of blazar heating
 - Thermal history of the IGM
 - The Lyman- α forest
- 3 Structure formation
 - Entropy evolution
 - Bimodality of galaxy clusters
 - Formation of dwarf galaxies



Outline

- 1 Physics of blazar heating
 - TeV emission from blazars
 - Propagation of TeV photons
 - Plasma instabilities
- 2 The intergalactic medium
 - Properties of blazar heating
 - Thermal history of the IGM
 - The Lyman- α forest
- 3 Structure formation
 - Entropy evolution
 - Bimodality of galaxy clusters
 - Formation of dwarf galaxies




TeV gamma-ray astronomy

H.E.S.S.



MAGIC I



VERITAS



MAGIC II




The TeV gamma-ray sky

There are several classes of TeV sources:

- Galactic - pulsars, BH binaries, supernova remnants
- Extragalactic - **mostly** blazars, two starburst galaxies


VHE γ -ray Sky Map
($E_{\gamma} > 100$ GeV)




2011-01-08 - Up-to-date plot available at <http://www.inpp.mpg.de/~wagner/sources/>



Unified model of active galactic nuclei



The blazar sequence



Ghisellini (2011)

Propagation of TeV photons

- 1 TeV photons can pair produce with 1 eV photons:

$$\gamma + \gamma \rightarrow e^+ + e^-$$

- mean free path for this depends on the density of 1 eV photons:
 - typically ~ 100 Mpc
 - pairs produced with energy of 0.5 TeV ($\gamma = 10^6$)
- these pairs inverse Compton scatter off the CMB photons
 - mean free path is ~ 30 kpc
 - producing gamma-rays of ~ 1 GeV


$$E \sim \gamma^2 E_{\text{CMB}} \sim 1 \text{ GeV}$$

- each TeV point source should also be a GeV point source



What about the cascade emission?

Every TeV source should be associated with a 1-100 GeV gamma-ray halo – **not seen!**




Neronov & Vovk (2010)



Missing plasma physics?

How do beams of e^+/e^- propagate through the IGM?

- plasma processes are important
- interpenetrating beams of charged particles are unstable
- consider the two-stream instability:



- one frequency (timescale) and one length in the problem:

$$\frac{\omega_p}{\gamma} = \sqrt{\frac{4\pi e^2 n_e}{\gamma^2 m_e}}$$


$$\lambda_p = \frac{\gamma c}{\omega_p}$$



Two-stream instability: mechanism

wave-like perturbation with $\mathbf{k} \parallel \mathbf{v}_{\text{beam}}$, longitudinal charge oscillations in background plasma (Langmuir wave):


- initially homogeneous beam- e^- :
attractive (repulsive) force by potential maxima (minima)
- e^- attain lowest velocity in potential minima \rightarrow bunching up
- e^+ attain lowest velocity in potential maxima \rightarrow bunching up




Two-stream instability: mechanism

wave-like perturbation with $\mathbf{k} \parallel \mathbf{v}_{\text{beam}}$, longitudinal charge oscillations in background plasma (Langmuir wave):

- beam- e^+/e^- couple in phase with the background perturbation: enhances background potential
- stronger forces on beam- $e^+/e^- \rightarrow$ positive feedback
- exponential wave-growth \rightarrow instability



Two-stream instability: energy transfer





- energy is transferred to the plasma wave from particles with $v \gtrsim v_{phase} \rightarrow$ growing modes
- energy is transferred from the plasma wave to particles with $v \lesssim v_{phase} \rightarrow$ damped modes



Oblique instability

k oblique to \mathbf{v}_{beam} : real world perturbations don't choose "easy" alignment = \sum all orientations




Bret (2009), Bret+ (2010)



Beam physics – growth rates

excluded for collective
 plasma phenomena



- consider a light beam penetrating into relatively dense plasma

- maximum growth rate

$$\sim 0.4 \gamma \frac{n_{\text{beam}}}{n_{\text{IGM}}} \omega_p$$

- oblique instability beats IC by two orders of magnitude

Broderick, Chang, C.P. (2011)



Beam physics – growth rates

- non-linear evolution of these instabilities at these density contrasts is not known
- expectation from PIC simulations suggest substantial isotropization of the beam
- plasma instabilities dissipate the beam's energy, no energy left over for inverse Compton scattering off the CMB



Summary: Heating by TeV blazars

- blazars emit TeV gamma-rays
- production of e^+ / e^- -pairs with extragalactic-background-light photons
- energy of e^+ / e^- -pairs is dissipated locally by plasma instabilities \rightarrow heating the IGM
- heating is almost independent of density for $z < 3.5$ (underdense regions receive more energy per unit mass)




Outline

- 1 Physics of blazar heating
 - TeV emission from blazars
 - Propagation of TeV photons
 - Plasma instabilities
- 2 The intergalactic medium
 - Properties of blazar heating
 - Thermal history of the IGM
 - The Lyman- α forest
- 3 Structure formation
 - Entropy evolution
 - Bimodality of galaxy clusters
 - Formation of dwarf galaxies



TeV blazar luminosity density




Broderick, Chang, C.P. (2011)

- collect luminosity of all 23 TeV blazars with good spectral measurements
- account for the selection effects
- TeV blazar luminosity density is a scaled version ($\sim 0.2\%$) of that of quasars!
- assume that they trace each other for all z



Evolution of the heating rates



Chang, Broderick, C.P. (2011)



Blazar heating vs. photoheating

- total power from AGN/stars vastly exceeds the TeV power of blazars
- $T_{\text{IGM}} \sim 10^4$ K (1 eV) at mean density ($z \sim 2$)

$$\varepsilon_{\text{th}} = \frac{kT}{m_p c^2} \sim 10^{-9}$$

- radiative energy ratio emitted by BHs in the Universe (Fukugita & Peebles 2004)

$$\varepsilon_{\text{rad}} = \eta \Omega_{\text{bh}} \sim 0.1 \times 10^{-4} \sim 10^{-5}$$

- fraction of the energy energetic enough to ionize H I is ~ 0.1 :


$$\varepsilon_{\text{UV}} \sim 0.1 \varepsilon_{\text{rad}} \sim 10^{-6} \quad \rightarrow \quad kT \sim \text{keV}$$

- photoheating efficiency $\eta_{\text{ph}} \sim 10^{-3} \quad \rightarrow \quad kT \sim \eta_{\text{ph}} \varepsilon_{\text{UV}} m_p c^2 \sim \text{eV}$
 (limited by the abundance of H I/He II due to the small recombination rate)

- blazar heating efficiency $\eta_{\text{bh}} \sim 10^{-3} \quad \rightarrow \quad kT \sim \eta_{\text{bh}} \varepsilon_{\text{rad}} m_p c^2 \sim 10 \text{ eV}$
 (limited by the total power of TeV sources)



Thermal history of the IGM




Chang, Broderick, C.P. (2011)




Evolution of the temperature-density relation

no blazar heating



blazar heating




Chang, Broderick, C.P. (2011)

- blazars and extragalactic background light are uniform
→ blazar heating independent of density
→ causes inverted temperature-density relation, $T \propto 1/\delta$
- blazars completely change the thermal history of the diffuse IGM and late-time structure formation




Evolution of the temperature-density relation

no blazar heating



blazar heating



Chang, Broderick, C.P. (2011)

- blazars and extragalactic background light are uniform
→ blazar heating independent of density
→ causes inverted temperature-density relation, $T \propto 1/\delta$
- blazars completely change the thermal history of the diffuse IGM and late-time structure formation




Simulations with blazar heating

Puchwein, C.P., Springel, Broderick, Chang (2011):

- $L = 15h^{-1}\text{Mpc}$ boxes with 2×384^3 particles
- one reference run without blazar heating
- three with blazar heating at different levels of efficiency (to account for uncertainties in the expected blazar-heating rate)
- used an up-to-date model of the UV background (Faucher-Giguère et al. 2009)




Temperature-density relation



Puchwein, C.P., Springel, Broderick, Chang (2011)





Ly- α spectra



Puchwein+ (2011)




Ly- α flux PDFs and power spectra



Puchwein+ (2011)


Voigt profile decomposition



- decomposing Lyman- α forest into individual Voigt profiles
- allows studying the thermal broadening of absorption lines



Voigt profile decomposition – line width distribution



Puchwein+ (2011)



Lyman- α forest in a blazar heated Universe

impressive improvement in modelling the Lyman- α forest is a direct consequence of the peculiar properties of blazar heating:

- **heating rate independent of IGM density** \rightarrow naturally produces the inverted $T-\rho$ relation that Lyman- α forest data demand
- **recent and continuous nature of the heating** needed to match the redshift evolutions of all Lyman- α forest statistics
- **magnitude of the heating rate required by Lyman- α forest data** \sim the total energy output of TeV blazars (or equivalently $\sim 0.2\%$ of that of quasars)




Outline

- 1 Physics of blazar heating
 - TeV emission from blazars
 - Propagation of TeV photons
 - Plasma instabilities
- 2 The intergalactic medium
 - Properties of blazar heating
 - Thermal history of the IGM
 - The Lyman- α forest
- 3 **Structure formation**
 - Entropy evolution
 - Bimodality of galaxy clusters
 - Formation of dwarf galaxies




Entropy evolution

temperature evolution



entropy evolution




C.P., Chang, Broderick (2011)

- evolution of the entropy, $K_e = kTn_e^{-2/3}$, at mean density
- blazar heating substantially increases the entropy floor ($z \lesssim 2$)




Evolution of the entropy-density relation

no blazar heating



blazar heating



C.P., Chang, Broderick (2011)

- blazar heating substantially increases the entropy in voids
- scatter is also increased → larger stochasticity of structure formation



Blazar heating: AGN feedback vs. pre-heating


Blazar heating is an amalgam of pre-heating and AGN feedback:

- **blazar heating is not localized** (\neq AGN feedback)
→ may change initial conditions for forming groups (but provides no stability for cool cores, CCs)
- **blazar heating generates time-dependent entropy floor** (\neq pre-heating)
→ may solve the classical problems of pre-heating ($z \sim 3$):
 - provides a physical mechanism
 - does not starve galaxy formation for $z \lesssim 3$
 - early forming groups can cool and develop observed low- K_e cores




Mass accretion history of groups/clusters

mass accretion history



mass accretion rates




C.P., Chang, Broderick (2011)

- peak entropy injection from blazar heating ($z \sim 1$) matches formation time of groups
- early forming groups are unaffected and develop cool cores
- late forming groups may have an elevated entropy core




Entropy profiles: effect of blazar heating

varying formation time



varying cluster mass



C.P., Chang, Broderick (2011)

If significant fraction of intra-group medium collapses from IGM:

- z-dependent excess entropy in cores (no cooling)
- largest effect for late forming, small objects




Scenario for the bimodality of cluster core entropies?

- entropy core, $K_{e,0}$, immediately after formation is set by the z -dependent blazar heating
- only late forming groups ($z \lesssim 1$) are directly affected by blazar (pre-)heating
- if the cooling time, t_{cool} , is shorter than the time period to the successive merger, t_{merger} , the group will radiate away the elevated core entropy and evolve into a CC
- if $t_{\text{cool}} > t_{\text{merger}}$, merger shocks can gravitationally reprocess the entropy cores and amplify them \rightarrow potentially those forming clusters evolve into non-cool core (NCC) systems



Gravitational reprocessing of entropy floors




Borgani+ (2005)

- larger $K_{e,0}$ of a merging cluster facilitates shock heating \rightarrow increase of $K_{e,0}$ over entropy floor
- entropy floor of 100 keV cm^2 at $z = 3$ in non-radiative simulation:
net entropy amplification factor $\sim 3\text{--}5$ for clusters and groups (Borgani+ 2005)
- expect median of $K_{e,0} \sim 150 \text{ keV cm}^2$; maximum $K_{e,0} \sim 600 \text{ keV cm}^2$



Bimodality of cluster core entropies



Cavagnolo+ (2009)

- *Chandra* observations match blazar heating expectations!
- need hydrodynamic simulations to confirm this scenario



Jeans mass

- on small enough scales, the thermal pressure can oppose gravitational collapse of the gas
- characteristic length scale below which objects will not form
- Jeans wavenumber and mass is obtained by balancing the sound crossing and free-fall timescales

$$k_J(a) \equiv \frac{a}{c_s(a)} \sqrt{4\pi G \bar{\rho}(a)}$$


$$M_J(a) \equiv \frac{4\pi}{3} \bar{\rho}(a) \left(\frac{2\pi a}{k_J(a)} \right)^3 = \frac{4\pi^{5/2}}{3} \frac{c_s^3(a)}{G^{3/2} \bar{\rho}^{1/2}(a)}$$

- blazar heating increases the IGM temperature by ~ 10 :

$$\frac{M_{J,\text{blazar}}}{M_{J,\text{photo}}} = \left(\frac{c_{s,\text{blazar}}}{c_{s,\text{photo}}} \right)^3 = \left(\frac{T_{\text{blazar}}}{T_{\text{photo}}} \right)^{3/2} \gtrsim 30$$



Filtering mass – dwarf formation




C.P., Chang, Broderick (2011)




Peebles' void phenomenon explained?

mean density



void, $1 + \delta = 0.5$




C.P., Chang, Broderick (2011)

- blazar heating efficiently suppresses the formation of void dwarfs within existing DM halos of masses $< 3 \times 10^{11} M_\odot$ ($z = 0$)
- reconciling the number of void dwarfs in simulations and the paucity of those in observations




“Missing satellite” problem in the Milky Way

satellite formation time



satellite luminosity function



- blazar heating suppresses late satellite formation, reconciling low observed dwarf abundances with CDM simulations




Conclusions on blazar heating

- **novel mechanism; dramatically alters thermal history of the IGM:**
 - uniform and z -dependent preheating
 - rate independent of density \rightarrow inverted $T-\rho$ relation
 - consistent picture of Lyman- α forest
- **significantly modifies late-time structure formation:**
 - group/cluster bimodality of core entropy values
 - may suppress Sunyaev-Zel'dovich power spectrum
 - dwarf formation: “missing satellite” problem, void phenomenon
- **explains puzzles in high-energy astrophysics:**
 - TeV blazars can evolve like quasars
 - extragalactic gamma-ray background at $E \gtrsim 10$ GeV
 - invalidates intergalactic B -constraints from blazar spectra



How efficient is heating by AGN feedback?



C.P., Chang, Broderick (2011)

- on a buoyancy timescale, no AGN outburst transforms a CC to a non-cool core (NCC) cluster!

- cavity enthalpy

$$E_{\text{cav}} = 4 PV_{\text{tot}}$$

- in some cases

$$E_{\text{cav}} \gtrsim E_{\text{bind}}(R_{2500})$$

- cavity energy only couples weakly into ICM, but prevents cooling catastrophe

